

Table 1. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trapper, and percent successful trappers by furbearer species for the 2018-19 season. There were 782 trappers who purchased a license for the 2018-19 season. Correcting for the percentage of license holders who indicated they did not trap or trapped feral hogs only (28.2%), there was an estimated N=561 licensed furbearer trappers in Mississippi in 2018-19 to which estimates were extrapolated.

Species	Harvest ^a	SE	Average Catch Per Trapper ^a	SE	Average Catch Per Successful Trapper	SE	N ^b	Percent Successful Trappers ^a	SE
Mink	32	6.87	0.06	0.01	2.50	0.41	10	2.29	0.34
Raccoon	9,020	318.56	16.08	0.57	20.37	0.68	345	78.95	0.92
Muskrat	312	48.17	0.56	0.09	8.68	1.13	28	6.41	0.55
Red Fox	193	13.55	0.34	0.02	1.97	0.10	76	17.39	0.85
Gray Fox	553	29.98	0.99	0.05	3.48	0.14	124	28.38	1.02
Bobcat	691	34.78	1.23	0.06	3.76	0.15	143	32.72	1.06
Opossum	5,522	221.22	9.84	0.39	16.48	0.59	261	59.73	1.10
Otter	959	71.48	1.71	0.13	6.33	0.41	118	27.00	1.00
Spotted Skunk	9	3.13	0.02	0.01	2.33	0.63	3	0.69	0.19
Striped Skunk	281	27.26	0.50	0.05	3.65	0.29	60	13.73	0.77
Coyote	2,677	129.33	4.77	0.23	10.43	0.44	200	45.77	1.12
Weasel	0	0.00	0.00	0.00	0.00	0.00	0	0.00	0.00
Nutria	1,018	200.57	1.81	0.28	13.00	1.89	61	13.96	0.78
Beaver	5,501	327.05	9.81	0.58	19.57	1.08	219	50.11	1.13

^a n = 437

^b n=sample size for average catches per successful trapper