

Table 1. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trapper, and percent successful trappers by furbearer species for the 1976-77 season.

Species	Harvest ^a	SE	Catch Per Trapper ^a	SE	Average Catch Per Successful Trapper	SE	n ^b	Percent Successful Trappers ^a	SE
Mink	22,105	1,566	7.67	0.54	11.80	0.78	403	65.0	1.7
Raccoon	61,528	3,247	21.36	1.13	25.61	1.29	517	83.4	1.3
Muskrat	41,473	3,686	14.40	1.28	31.21	2.50	286	46.1	1.8
Red Fox	6,301	506	2.19	0.18	4.97	0.35	273	44.0	1.8
Gray Fox	6,454	1,135	2.24	0.39	19.56	2.86	71	11.5	1.1
Bobcat	4,517	429	1.57	0.15	4.54	0.37	214	34.5	1.7
Opossum	49,098	2,986	17.04	1.04	24.63	1.38	429	69.2	1.6
Otter	1,482	233	0.51	0.08	4.31	0.54	74	11.9	1.2
Spotted Skunk	1,915	305	0.66	0.11	5.64	0.72	73	11.8	1.1
Striped Skunk	2,537	321	0.88	0.11	5.15	0.51	106	17.1	1.3
Coyote	469	91	0.16	0.03	2.53	0.35	40	6.5	0.9
Weasel	274	127	0.10	0.04	2.81	1.21	21	3.4	0.6
Nutria	4,308	1,105	1.50	0.38	13.06	3.11	71	11.5	1.1
Beaver	19,326	2,087	6.71	0.72	16.00	1.59	260	41.9	1.8

^a n = 620

^b - Sample size for average catches per successful trapper

Table 2. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trapper, and percent successful trappers by furbearer species for the 1977-78 season.

Species	Harvest ^a	SE	Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	15,791	1,730	6.68	0.73	10.80	1.12	281	61.9	2.1
Raccoon	52,780	3,557	22.29	1.47	24.44	1.57	415	91.4	1.2
Muskrat	29,772	4,229	12.60	1.78	28.19	3.76	203	44.7	2.1
Red Fox	4,090	429	1.73	0.18	4.18	0.38	188	41.4	2.1
Gray Fox	13,616	1,667	5.75	0.70	9.66	1.12	271	59.7	2.1
Bobcat	3,273	408	1.39	0.17	4.13	0.45	152	33.5	2.0
Opossum	36,972	2,379	15.60	0.98	20.42	1.18	348	76.7	1.8
Otter	973	177	0.41	0.07	3.07	0.45	61	13.4	1.4
Spotted Skunk	1,212	206	0.51	0.09	4.09	0.53	57	12.6	1.4
Striped Skunk	1,753	265	0.74	0.11	4.43	0.52	76	16.7	1.6
Coyote	395	112	0.17	0.05	2.62	0.62	29	6.4	1.0
Weasel	68	19	0.03	0.01	1.18	0.11	11	2.4	0.6
Nutria	5,692	2,369	2.41	1.00	24.31	9.73	45	9.9	1.3
Beaver	15,984	1,901	6.76	0.80	14.63	1.59	210	46.3	2.1

^a n = 527

^b n = 454

^c Sample size for average catches per successful trapper

Table 3. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trapper, and percent successful trappers by furbearer species for the 1978-79 season.

Species	Harvest ^a	SE	Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper ^b	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	12,879	566	4.97	0.22	7.99	0.32	683	62.4	1.1
Raccoon	59,200	1,808	22.90	0.70	24.67	0.74	1,017	92.9	0.6
Muskrat	20,354	1,325	7.83	0.51	18.01	1.07	479	43.7	1.1
Red Fox	4,667	264	1.81	0.10	3.86	0.20	512	46.8	1.1
Gray Fox	13,051	596	5.05	0.23	8.11	0.34	682	62.3	1.1
Bobcat	3,263	194	1.26	0.07	3.51	0.18	394	36.0	1.1
Opossum	52,310	1,608	20.24	0.62	23.43	0.69	946	86.4	0.8
Otter	1,045	86	0.39	0.03	2.72	0.17	163	14.9	0.8
Spotted Skunk	1,841	152	0.71	0.06	4.00	0.27	195	17.8	0.9
Striped Skunk	3,610	235	1.40	0.09	5.48	0.29	279	25.5	1.0
Coyote	779	102	0.27	0.04	2.77	0.32	119	10.9	0.7
Weasel	94	22	0.04	0.01	1.67	0.31	24	2.2	0.3
Nutria	3,164	485	1.22	0.19	12.19	1.68	110	10.0	0.6
Beaver	17,192	1,038	6.64	0.40	13.85	0.77	526	48.0	1.1

^a n = 1,104

^b n = 1,095

^c Sample size for average catches per successful trapper

Table 4. Expanded statewide harvest estimates, average catch per license holder, average catch per successful trapper, and percent successful license holders by furbearer species for the 1979-80 season.

Species	Harvest ^a	SE	Catch Per License Holder ^a	SE	Average Catch Per Successful Trapper	SE	n ^b	Percent Successful Trappers ^a	SE
Mink	20,259	1,310	5.49	0.35	8.87	0.55	738	61.9	1.0
Raccoon	72,053	2,141	19.52	0.58	20.94	0.61	1,112	93.2	0.6
Muskrat	38,899	3,050	10.54	0.83	24.75	1.82	508	42.6	1.2
Red Fox	4,415	380	1.20	0.10	3.53	0.28	404	33.9	1.1
Gray Fox	12,475	615	3.40	0.17	6.26	0.28	644	54.0	1.2
Bobcat	3,066	183	0.83	0.05	2.84	0.13	349	29.3	1.1
Opossum	58,088	1,875	15.74	0.51	18.72	0.57	1,003	84.1	0.9
Otter	1,532	145	0.41	0.04	2.80	0.21	177	14.8	0.8
Spotted Skunk	1,581	205	0.43	0.06	3.81	0.43	134	11.2	0.8
Striped Skunk	2,729	236	0.74	0.06	4.26	0.29	207	17.4	0.9
Coyote	1,151	148	0.31	0.04	2.60	0.29	143	12.0	0.8
Weasel	111	35	0.03	0.01	2.00	0.50	18	1.5	0.3
Nutria	7,821	1,513	2.12	0.41	19.30	3.51	131	11.0	0.7
Beaver	19,352	1,104	5.24	0.30	11.25	0.57	556	46.6	1.2

^a n = 1,193

^b - Sample size for average catches per successful trapper

Table 5. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trapper, and percent successful trappers by furbearer species for the 1980-81 season.

Species	Harvest ^a	SE	Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	16,754	771	5.60	0.26	8.66	0.38	897	65.8	0.9
Raccoon	49,594	1,182	16.67	0.39	18.26	0.41	1,259	92.4	0.5
Muskrat	30,185	1,728	10.23	0.59	21.11	1.12	663	48.6	1.0
Red Fox	2,487	146	0.84	0.05	2.83	0.14	408	29.9	0.9
Gray Fox	7,965	320	2.70	0.11	5.51	0.19	670	49.2	0.9
Bobcat	2,230	129	0.76	0.04	2.85	0.13	363	26.6	0.9
Opossum	27,235	848	9.19	0.28	13.08	0.37	965	70.8	0.9
Otter	1,044	80	0.35	0.03	2.56	0.15	189	13.9	0.7
Spotted Skunk	1,048	107	0.36	0.04	3.74	0.31	130	9.5	0.6
Striped Skunk	2,869	387	0.97	0.13	5.81	0.74	229	16.8	0.7
Coyote	947	70	0.32	0.02	2.35	0.13	187	13.7	0.7
Weasel	22	5	0.01	<0.01	1.00	0.00	10	0.7	0.2
Nutria	4,765	693	1.62	0.24	12.84	1.75	172	12.6	0.7
Beaver	18,162	926	6.15	0.31	12.33	0.58	683	50.01	1.0

^a n = 1,452

^b n = 1,363

^c Sample size for average catches per successful trapper

Table 6. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trapper, and percent successful trappers by furbearer species for the 1981-82 season.

Species	Harvest ^a	SE	Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	15,064	859	5.21	0.30	7.90	0.43	774	66.6	1.1
Raccoon	46,320	1,258	16.04	0.43	17.42	0.45	1,079	92.8	0.6
Muskrat	27,100	1,533	9.43	0.53	19.95	1.02	551	47.4	1.1
Red Fox	2,026	114	0.70	0.04	2.51	0.11	328	28.2	1.0
Gray Fox	5,936	267	2.06	0.09	4.32	0.16	557	47.9	1.1
Bobcat	1,942	124	0.68	0.04	2.83	0.14	278	23.9	1.0
Opossum	26,186	977	9.08	0.34	13.85	0.46	767	66.0	1.1
Otter	833	68	0.29	0.02	2.32	0.13	146	12.6	0.7
Spotted Skunk	887	130	0.31	0.05	4.24	0.52	85	7.3	0.5
Striped Skunk	2,477	213	0.86	0.07	5.15	0.36	195	16.8	0.8
Coyote	1,104	81	0.38	0.03	2.30	0.12	195	16.8	0.8
Weasel	44	12	0.02	<0.01	1.38	0.24	13	1.1	0.2
Nutria	6,692	1,482	2.33	0.52	16.55	3.56	164	14.1	0.7
Beaver	14,694	893	5.12	0.31	10.53	0.59	566	48.7	1.1

^a n = 1,228

^b n = 1,163

^c Sample size for average catches per successful trapper

Table 7. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trapper and percent successful trappers by furbearer species for the 1982-83 season.

Species	Harvest ^a	SE	Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	9,911	677	4.57	0.31	8.12	0.52	497	56.9	1.3
Raccoon	35,247	1,232	16.28	0.56	17.68	0.59	812	92.9	0.7
Muskrat	16,752	1,216	7.73	0.56	18.85	1.23	362	41.4	1.3
Red Fox	1,640	121	0.76	0.06	2.66	0.16	251	28.7	1.2
Gray Fox	5,090	288	2.37	0.13	4.67	0.23	444	50.8	1.3
Bobcat	1,270	87	0.58	0.04	2.43	0.12	213	24.4	1.1
Opossum	24,151	1,277	11.22	0.59	17.25	0.84	570	65.2	1.2
Otter	535	54	0.25	0.03	2.20	0.15	99	11.3	0.8
Spotted Skunk	651	127	0.30	0.06	3.63	0.64	73	8.4	0.7
Striped Skunk	1,923	163	0.89	0.08	4.96	0.32	158	18.1	1.0
Coyote	1,073	102	0.50	0.05	2.68	0.21	163	18.6	1.0
Weasel	10	4	<0.01	<0.01	1.00	0.00	4	0.5	0.2
Nutria	2,581	460	1.20	1.20	10.84	1.76	97	11.1	0.8
Beaver	8,236	688	3.79	3.79	9.56	0.73	351	40.2	1.3

^a n = 979

^b n = 874

^c Sample size for average catches per successful trapper

Table 8. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trapper and percent successful trappers by furbearer species for the 1983-84 season.

Species	Harvest ^a	SE	Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	8,776	683	5.61	0.44	9.36	0.69	395	60.1	1.5
Raccoon	27,427	1,041	17.59	1.65	19.53	0.70	592	90.1	0.9
Muskrat	15,749	2,065	10.05	1.32	23.37	2.96	284	43.2	1.5
Red Fox	1,165	94	0.74	0.06	2.65	0.17	185	28.2	1.3
Gray Fox	3,383	219	2.17	0.14	4.80	0.27	297	45.2	1.5
Bobcat	1,276	90	0.82	0.06	2.83	0.15	190	28.9	1.3
Opossum	15,619	734	10.02	0.46	15.98	0.64	412	62.7	1.4
Otter	444	41	0.28	0.03	2.05	0.11	91	13.9	1.0
Spotted Skunk	415	54	0.27	0.03	2.87	0.26	61	9.3	0.9
Striped Skunk	1,597	154	1.02	0.10	4.95	0.38	136	20.7	1.2
Coyote	1,108	101	0.71	0.06	2.99	0.22	156	23.7	1.3
Weasel	10	4	0.01	<0.01	1.00	0	4	0.6	0.2
Nutria	2,081	576	1.33	0.37	10.96	2.92	80	12.2	1.0
Beaver	6,033	592	3.87	0.38	10.34	0.93	246	37.4	1.4

^a n = 714

^b n = 657

^c Sample size for average catches per successful trapper

Table 9. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trapper and percent successful trappers by furbearer species for the 1984-85 season.

Species	Harvest ^a	SE	Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	8,682	572	5.20	0.34	8.51	0.52	419	61.2	1.4
Raccoon	38,696	1,302	23.19	0.77	24.41	0.79	651	95.0	0.6
Muskrat	17,266	2,269	10.34	1.36	25.05	3.17	283	41.3	1.4
Red Fox	1,500	141	0.90	0.08	3.29	0.27	187	27.3	1.3
Gray Fox	3,702	195	2.21	0.12	4.83	0.20	315	46.0	1.5
Bobcat	1,371	87	0.82	0.05	2.71	0.12	208	30.4	1.3
Opossum	15,573	870	9.34	0.52	15.41	0.78	415	60.6	1.4
Otter	648	53	0.39	0.03	2.02	0.11	132	19.3	1.2
Spotted Skunk	244	31	0.13	0.02	2.24	0.17	41	6.0	0.7
Striped Skunk	1,383	154	0.83	0.09	4.85	0.44	117	17.1	1.1
Coyote	1,680	150	1.01	0.09	3.65	0.28	189	27.6	1.3
Weasel	10	4	<0.01	<0.01	1.00	0.00	4	0.6	0.2
Nutria	2,292	459	1.37	0.28	11.76	2.17	80	11.7	0.9
Beaver	9,734	994	5.84	0.59	12.77	1.24	313	45.7	1.5

^a n = 710

^b n = 685

^c Sample size for average catches per successful trapper

Table 10. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trapper and percent successful trappers by furbearer species for the 1985-86 season.

Species	Harvest ^a	SE	Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	7,642	602	6.00	0.47	9.05	0.68	357	66.5	1.5
Raccoon	34,572	1,452	27.11	1.12	28.71	1.16	509	94.8	0.7
Muskrat	10,530	998	80.23	0.78	20.42	1.75	218	40.6	1.6
Red Fox	887	78	0.69	0.06	2.68	0.18	140	26.1	1.4
Gray Fox	2,468	165	1.92	0.13	4.46	0.24	234	43.6	1.6
Bobcat	1,254	92	0.98	0.07	2.85	0.16	186	34.6	1.6
Opossum	11,775	625	9.23	0.49	14.9	0.68	334	62.2	1.6
Otter	733	68	0.58	0.05	2.42	0.17	128	23.8	1.4
Spotted Skunk	263	47	0.20	0.04	3.83	0.44	29	5.4	0.7
Striped Skunk	1,261	144	0.99	0.11	5.03	0.47	106	19.7	1.3
Coyote	1,306	151	1.03	0.12	3.63	0.38	152	28.3	1.5
Weasel	7	3	0.01	<0.01	1.00	0	3	0.6	0.2
Nutria	9,355	5,426	7.35	4.27	54.16	31.28	73	13.6	1.1
Beaver	10,247	924	8.04	0.72	16.28	1.36	266	49.5	1.6

^a n = 574

^b n = 537

^c Sample size for average catches per successful trapper

Table 11. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers for furbearer species for the 1986-87 season.

Species	Harvest ^a	SE	Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	8,881	684	5.89	0.46	9.32	0.68	393	64	1.5
Raccoon	41,400	1,439	27.50	0.50	28.60	0.60	597	97.2	0.5
Muskrat	12,142	1,207	8.10	0.81	21.31	1.94	235	38.3	1.5
Red Fox	921	67	0.59	0.04	2.33	0.12	163	26.5	1.4
Gray Fox	4,192	247	2.70	0.15	5.29	0.27	327	53.3	1.5
Bobcat	1,945	138	1.29	0.09	3.41	0.20	235	38.3	1.5
Opossum	20,078	1,017	13.42	0.68	20.15	0.91	411	66.9	1.5
Otter	955	76	0.64	0.05	2.64	0.15	149	24.3	1.3
Spotted Skunk	347	63	0.23	0.04	3.76	1.51	38	6.2	0.7
Striped Skunk	1,523	150	1.02	0.10	5.06	0.39	124	20.2	1.2
Coyote	1,804	169	1.21	0.11	4.18	0.34	178	29.0	1.4
Weasel	15	6	0.01	<0.01	1.50	0.38	4	0.7	0.2
Nutria	4,199	810	2.80	0.54	15.46	2.81	112	18.2	1.2
Beaver	15,214	2,711	10.16	1.82	20.18	3.54	311	50.7	1.5

^a n = 643

^b n = 614

^c Sample size for average catches per successful trapper

Table 12. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 1987-88 season.

Species	Harvest ^a	SE	Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	6,201	689	3.90	0.43	7.46	0.78	246	52.2	1.9
Raccoon	41,898	2,058	26.32	1.26	27.68	1.31	448	95.1	0.8
Muskrat	9,097	1,445	5.72	0.91	19.51	2.87	138	29.3	1.8
Red Fox	1,115	107	0.70	0.07	2.68	0.19	123	26.1	1.7
Gray Fox	5,670	408	3.56	0.25	7.02	0.42	239	50.7	1.9
Bobcat	2,521	225	1.58	0.14	4.60	0.33	162	34.4	1.8
Opossum	16,632	1,164	10.43	0.72	17.15	1.06	287	60.9	1.9
Otter	1,064	106	0.67	0.07	3.09	0.20	102	21.7	1.6
Spotted Skunk	301	73	0.19	0.05	3.56	0.64	25	5.3	0.9
Striped Skunk	1,531	206	0.96	0.13	4.92	0.54	92	19.5	1.5
Coyote	2,707	354	1.68	0.22	5.52	0.65	145	30.8	1.8
Weasel	17	7	0.01		1.25	0.21	4	0.8	0.4
Nutria	1,602	269	1.01	<0.01 0.17	6.77	0.95	70	14.9	1.4
Beaver	12,952	1,623	8.01	1.01	17.5	2.06	219	46.5	1.9

^a n = 509

^b n = 471

^c Sample size for average catches per successful trapper

Table 13. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 1988-89 season.

Species	Harvest ^a	SE	Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	5,291	511	5.77	0.55	11.50	1.02	212	50.2	1.7
Raccoon	17,901	816	19.55	0.86	20.67	0.89	399	94.5	0.8
Muskrat	6,853	964	7.48	1.05	25.47	3.29	124	29.3	1.6
Red Fox	853	69	0.93	0.07	3.25	0.19	121	28.6	1.6
Gray Fox	3,980	272	4.35	0.29	7.77	0.46	236	55.9	1.8
Bobcat	1,686	122	1.84	0.13	4.32	0.25	180	42.4	1.8
Opossum	7,209	414	7.87	0.44	13.56	0.64	245	58.1	1.8
Otter	532	51	0.58	0.06	2.66	0.18	92	21.8	1.5
Spotted Skunk	243	52	0.19	0.05	4.48	0.73	25	5.9	0.8
Striped Skunk	788	87	0.86	0.09	4.37	0.36	93	19.6	1.4
Coyote	2,941	369	3.21	0.40	7.92	0.93	171	40.5	1.8
Weasel	9	3	0.01	<0.01	1.00	0	4	0.9	0.3
Nutria	1,515	272	1.65	0.30	11.44	1.81	61	14.5	1.3
Beaver	9,753	996	10.65	1.08	19.89	1.91	226	53.6	1.8

^a n = 482

^b n = 422

^c Sample size for average catches per successful trapper

Table 14. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 1989-90 season.

Species	Harvest ^a	SE	Catch Per Trapper ^a	SE	Average Catch Per Successful Trapper	SE	n ^b	Percent Successful Trappers	SE
Mink	2,106	287	4.11	0.56	9.98	1.20	88	41.7	2.6
Raccoon	7,893	602	15.53	0.60	18.08	1.29	182	86.3	1.8
Muskrat	2,506	482	4.85	0.95	15.83	2.78	66	31.3	2.4
Red Fox	269	43	0.53	0.08	2.38	0.30	47	22.3	2.2
Gray Fox	1,511	147	2.99	0.29	6.84	0.49	101	47.9	2.6
Bobcat	568	68	7.81	0.13	3.29	0.31	72	34.1	2.4
Opossum	3,969	504	1.12	0.99	16.71	1.88	99	46.9	2.6
Otter	425	73	0.84	0.14	4.21	0.57	42	19.9	2.1
Spotted Skunk	79	39	0.16	0.08	4.71	2.01	7	3.3	0.9
Striped Skunk	307	57	0.60	0.11	5.12	0.59	25	11.8	1.7
Coyote	1,535	256	3.03	0.50	8.10	1.23	79	37.4	2.6
Weasel	0	0	0	0	0	0	0	0	0
Nutria	1,017	259	2.01	0.51	15.14	3.33	28	13.3	1.8
Beaver	8,519	1,610	16.14	3.12	33.48	6.01	107	50.7	2.6

^a n = 236

^b - Sample size for average catches per successful trapper

Table 15. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 1990-91 season.

Species	Harvest ^a	SE	Catch Per Trapper ^a	SE	Average Catch Per Successful Trapper	SE	n ^b	Percent Successful Trappers	SE
Mink	917	145	3.38	0.53	8.51	1.11	47	40.1	3.4
Raccoon	4,232	411	15.60	1.45	20.97	1.74	88	75.2	3.0
Muskrat	681	153	2.41	0.56	9.58	1.86	31	26.5	3.1
Red Fox	236	47	0.88	0.17	3.81	0.58	27	23.1	2.9
Gray Fox	839	114	3.08	0.41	6.78	0.74	54	46.2	3.5
Bobcat	362	60	1.32	0.22	4.51	0.57	35	29.9	3.2
Opossum	2,008	226	7.21	0.80	13.69	1.20	64	54.7	3.5
Otter	199	38	0.69	0.14	3.22	0.46	27	23.1	2.9
Spotted Skunk	46	23	0.17	0.09	6.67	2.18	3	2.6	1.1
Striped Skunk	131	32	0.49	0.12	4.07	0.65	14	12.0	2.2
Coyote	1,146	222	4.26	0.82	10.20	1.78	49	41.9	3.4
Weasel	0	0	0	0	0	0	0	0	0
Nutria	378	121	1.41	0.45	9.71	2.68	17	14.5	2.5
Beaver	4,475	872	15.93	3.20	33.66	6.06	58	49.6	3.5

^a n = 147

^b - Sample size for average catches per successful trapper

Table 16. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 1991-92 season.

Species	Harvest ^a	SE	Catch Per Trapper ^a	SE	Average Catch Per Successful Trapper	SE	n ^b	Percent Successful Trappers	SE
Mink	1,424	213	5.09	0.75	11.03	1.44	59	46.1	3.3
Raccoon	5,443	505	19.44	1.75	23.70	2.01	105	82.0	2.5
Muskrat	1,256	241	4.48	0.85	15.11	2.47	38	29.7	3.0
Red Fox	199	40	0.71	0.14	3.96	0.59	23	18.0	0.5
Gray Fox	1,092	187	3.90	0.66	9.78	1.48	51	39.8	3.2
Bobcat	360	64	1.29	0.23	4.53	0.65	36	28.1	2.9
Opossum	2,217	270	7.92	0.95	12.28	1.49	71	55.5	3.2
Otter	302	54	1.08	0.19	3.73	0.55	37	28.9	3.0
Spotted Skunk	9	4	0.03	0.01	1.33	0.25	3	2.3	1.0
Striped Skunk	291	64	1.03	0.23	5.54	0.98	24	18.8	2.6
Coyote	1,584	269	5.58	0.95	15.40	2.25	47	36.7	3.2
Weasel	0	0	0	0	0	0	0	0	0
Nutria	254	66	0.09	0.23	5.04	1.11	23	18.0	2.5
Beaver	5,110	1,319	18.25	4.70	32.44	8.16	72	56.3	3.2

^a n = 144

^b - Sample size for average catches per successful trapper

Table 17. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 1992-93 season.

Species	Harvest ^a	SE	Catch Per Trapper ^a	SE	Average Catch Per Successful Trapper	SE	n ^b	Percent Successful Trappers	SE
Mink	1,503	413	5.51	1.51	14.25	3.74	48	38.7	3.2
Raccoon	5,763	622	21.15	2.23	28.51	2.79	92	74.2	2.9
Muskrat	1,439	298	5.28	1.09	15.98	2.94	41	33.1	3.1
Red Fox	167	31	0.61	0.11	3.62	0.40	21	16.9	2.5
Gray Fox	1,050	179	3.85	0.65	9.02	1.36	53	42.7	3.3
Bobcat	483	73	1.77	0.27	5.24	0.19	42	33.9	3.2
Opossum	2,366	309	8.69	1.12	15.61	1.79	69	55.6	3.3
Otter	349	58	1.28	0.21	4.68	0.58	34	27.4	3.0
Spotted Skunk	26	8	0.09	0.03	1.50	0.28	8	6.5	1.6
Striped Skunk	426	81	1.56	0.29	6.47	0.96	30	24.2	2.9
Coyote	1,224	219	4.49	0.80	12.38	1.92	45	36.3	3.2
Weasel	0	0	0	0	0	0	0	0	0
Nutria	694	141	2.54	0.51	9.58	1.63	33	26.6	2.9
Beaver	6,203	1,264	22.77	4.61	41.51	8.05	68	54.8	N/A

^a n = 142

^b - Sample size for average catches per successful trapper

Table 18. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 1993-94 season.

Species	Harvest ^a	SE	Catch Per Trapper ^a	SE	Average Catch Per Successful Trapper	SE	n ^b	Percent Successful Trappers	SE
Mink	685	321	3.32	1.56	13.00	5.93	23	25.6	3.5
Raccoon	4,887	472	23.71	2.21	30.06	2.52	71	78.9	3.2
Muskrat	725	235	3.52	1.14	13.78	4.11	23	25.6	3.5
Red Fox	117	33	0.57	0.16	2.83	0.66	18	20.0	3.2
Gray Fox	643	102	3.12	0.49	7.39	0.94	38	42.2	3.9
Bobcat	474	80	2.30	0.38	5.45	0.76	38	42.2	3.9
Opossum	1,649	213	8.00	1.01	14.40	1.52	50	55.6	4.0
Otter	49	68	2.18	0.32	4.78	0.58	41	45.6	4.0
Spotted Skunk	44	22	0.21	0.11	4.75	1.87	4	4.4	1.6
Striped Skunk	336	135	1.78	0.65	13.33	4.24	12	13.3	2.7
Coyote	909	154	4.41	0.74	10.18	1.44	39	43.3	3.9
Weasel	5	2	0.002	0.01	1.00	0	2	2.2	1.2
Nutria	6,153	3,952	29.86	19.20	83.97	53.72	32	35.6	3.8
Beaver	6,563	1,249	31.84	3.01	50.28	9.02	57	63.3	3.8

^a n = 100

^b - Sample size for average catches per successful trapper

Table 19. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 1994-95 season.

Species	Harvest ^a	SE	Catch Per Trapper ^a	SE	Average Catch Per Successful Trapper	SE	n ^b	Percent Successful Trappers	SE
Mink	326	111	1.48	0.50	5.54	1.75	24	26.7	3.6
Raccoon	7,971	958	36.17	4.27	45.21	5.01	72	80.0	3.3
Muskrat	238	71	1.08	0.32	4.47	1.58	15	16.7	3.0
Red Fox	113	24	0.51	0.11	2.88	0.35	16	17.8	3.1
Gray Fox	531	82	2.41	0.37	6.03	0.70	36	40.0	4.0
Bobcat	419	80	1.90	0.36	4.75	0.77	36	40.0	4.0
Opossum	2,552	365	11.58	1.63	19.66	2.44	53	58.9	4.0
Otter	598	107	2.71	0.49	5.30	0.85	46	51.1	4.0
Spotted Skunk	12	8	0.06	0.04	2.50	1.15	2	2.2	1.2
Striped Skunk	88	25	0.40	0.12	3.27	0.64	44	12.2	2.7
Coyote	1,283	310	5.82	1.40	14.97	3.29	35	38.9	4.0
Weasel	0	0	0	0	0	0	0	0	0
Nutria	541	115	2.46	0.52	7.89	1.39	28	31.1	3.8
Beaver	5,929	1,317	26.90	5.94	37.83	8.14	64	71.1	3.7

^a n = 98

^b - Sample size for average catches per successful trapper

Table 20. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 1995-96 season.

Species	Harvest ^a	SE	Catch Per Trapper ^a	SE	Average Catch Per Successful Trapper	SE	n ^b	Percent Successful Trappers	SE
Mink	315	89	1.35	0.52	7.39	1.77	18	26.1	4.0
Raccoon	5,399	637	26.75	3.63	37.43	3.98	61	88.4	2.9
Muskrat	267	66	1.38	0.38	6.65	1.24	17	24.6	4.0
Red Fox	76	22	0.45	0.13	2.91	0.57	11	15.9	3.4
Gray Fox	400	67	2.14	0.39	5.63	0.70	30	43.5	4.6
Bobcat	293	51	1.71	0.30	4.28	0.56	29	42.0	4.5
Opossum	1,809	305	8.78	1.72	18.21	2.68	42	60.9	4.5
Otter	452	78	2.43	0.45	4.78	0.71	40	58.0	4.5
Spotted Skunk	38	21	0.23	0.13	8.00	1.52	2	2.9	1.5
Striped Skunk	97	23	0.49	0.13	2.93	0.45	14	20.3	27.7
Coyote	844	169	4.93	1.02	11.16	1.91	32	46.4	55.6
Weasel	0	0	0	0	0	0	0	0	0
Nutria	20	117	3.09	0.68	10.08	1.46	26	37.7	46.6
Beaver	5,654	956	31.42	5.74	43.47	6.88	56	79.7	87.7

^a n = 85

^b - Sample size for average catches per successful trapper

Table 21. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 1996-97 season.

Species	Harvest ^a	SE	Catch Per Trapper ^a	SE	Average Catch Per Successful Trapper	SE	n ^b	Percent Successful Trappers	SE
Mink	600	111	1.56	0.29	5.13	0.85	55	30.7	2.5
Raccoon	13,704	1,024	35.98	2.63	38.57	2.76	167	93.3	1.4
Muskrat	1,432	572	3.76	1.50	15.65	6.12	43	24.0	2.3
Red Fox	209	41	0.58	0.11	3.12	0.46	33	18.3	2.1
Gray Fox	1,323	199	3.47	0.52	7.78	1.07	80	44.7	2.7
Bobcat	885	94	2.32	0.24	5.07	0.44	82	45.8	2.7
Opossum	3,872	304	10.17	0.78	14.80	1.00	123	68.7	2.5
Otter	1,404	162	3.68	0.42	6.95	0.71	95	53.1	2.7
Spotted Skunk	72	39	0.19	0.10	5.67	2.82	6	3.4	1.0
Striped Skunk	189	35	0.50	0.09	3.07	0.42	29	16.2	2.0
Coyote	1,870	313	4.91	0.82	11.42	1.77	77	43.0	2.7
Weasel	4	2	0.01	0.01	1.00	N/A	2	1.1	0.6
Nutria	890	143	2.63	0.37	8.13	1.09	52	29.1	2.5
Beaver	11,404	1,390	29.58	3.62	46.13	5.27	116	64.8	2.6

^a n = 196

^b - Sample size for average catches per successful trapper

Table 22. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 1997-98 season.

Species	Harvest ^a	SE	Catch Per Trapper ^a	SE	Average Catch Per Successful Trapper	SE	n	Percent Successful Trappers	SE
Mink	674	N/A	1.94	N/A	8.28	N/A	N/A	23.39	N/A
Raccoon	11,610	N/A	33.30	N/A	38.27	N/A	N/A	87.14	N/A
Muskrat	759	N/A	2.17	N/A	10.08	N/A	N/A	21.64	N/A
Red Fox	250	N/A	0.72	N/A	3.62	N/A	N/A	19.88	N/A
Gray Fox	745	N/A	2.13	N/A	6.10	N/A	N/A	35.04	N/A
Bobcat	631	N/A	1.81	N/A	3.92	N/A	N/A	46.20	N/A
Opossum	3,378	N/A	9.69	N/A	18.23	N/A	N/A	53.22	N/A
Otter	922	N/A	2.61	N/A	4.87	N/A	N/A	54.39	N/A
Spotted Skunk	22	N/A	0.06	N/A	2.75	N/A	N/A	2.34	N/A
Striped Skunk	181	N/A	0.52	N/A	3.71	N/A	N/A	14.03	N/A
Coyote	1,509	N/A	4.25	N/A	11.06	N/A	N/A	39.18	N/A
Weasel	2	N/A	0.01	N/A	1.00	N/A	N/A	0.59	N/A
Nutria	1,004	N/A	2.87	N/A	10.06	N/A	N/A	28.66	N/A
Beaver	8,894	N/A	23.89	N/A	35.80	N/A	N/A	71.35	N/A

^a n = 194

Table 23. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 1998-99 season.

Species	Harvest ^a	SE	Catch Per Trapper ^a	SE	Average Catch Per Successful Trapper	SE	n	Percent Successful Trappers	SE
Mink	528	N/A	2.06	N/A	7.66	N/A	N/A	27.31	N/A
Raccoon	10,559	N/A	37.42	N/A	47.93	N/A	N/A	96.12	N/A
Muskrat	443	N/A	1.53	N/A	8.92	N/A	N/A	20.16	N/A
Red Fox	182	N/A	0.51	N/A	3.34	N/A	N/A	2.17	N/A
Gray Fox	519	N/A	1.57	N/A	4.58	N/A	N/A	45.74	N/A
Bobcat	519	N/A	1.81	N/A	4.41	N/A	N/A	43.41	N/A
Opossum	3,304	N/A	11.12	N/A	23.49	N/A	N/A	58.14	N/A
Otter	1,003	N/A	3.21	N/A	7.01	N/A	N/A	58.14	N/A
Spotted Skunk	186	N/A	0.72	N/A	13.71	N/A	N/A	5.42	N/A
Striped Skunk	184	N/A	0.50	N/A	4.75	N/A	N/A	15.50	N/A
Coyote	1,643	N/A	5.48	N/A	12.48	N/A	N/A	53.49	N/A
Weasel	0	N/A	0	N/A	0	N/A	N/A	0	N/A
Nutria	1,411	N/A	4.85	N/A	15.47	N/A	N/A	36.43	N/A
Beaver	9,620	N/A	32.81	N/A	45.96	N/A	N/A	85.27	N/A

^a n = 171

Table 24. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 1999-2000 season.

Species	Harvest ^a	SE	Catch Per Trapper ^a	SE	Average Catch Per Successful Trapper	SE	n ^b	Percent Successful Trappers	SE
Mink	511	137.37	1.84	0.49	1.11	2.75	27	16.56	0.02
Raccoon	8,596	708.58	30.98	2.49	39.15	2.98	129	79.14	0.02
Muskrat	819	348.32	2.95	1.25	17.81	7.42	27	16.56	0.02
Red Fox	213	39.49	0.77	0.14	3.13	0.51	40	24.54	0.02
Gray Fox	698	77.74	2.52	0.27	7.07	0.61	58	35.58	0.02
Bobcat	815	102.51	2.94	0.37	6.47	0.72	74	45.40	0.03
Opossum	3,602	468.31	12.98	1.67	23.51	2.84	90	55.22	0.03
Otter	1,166	104.84	4.20	0.37	8.56	0.61	80	49.08	0.03
Spotted Skunk	172	83.76	0.62	0.30	14.43	6.60	7	4.29	0.01
Striped Skunk	407	59.87	1.47	0.21	6.64	0.74	36	22.09	0.02
Coyote	2,213	330.11	7.98	1.18	17.12	2.37	76	46.63	0.03
Weasel	2	1.09	0.01	0	1.00	N/A	1	0.61	0.00
Nutria	509	70.25	1.88	0.25	7.12	0.77	42	25.77	0.02
Beaver	7,522	779.20	27.11	2.76	36.83	3.95	120	73.62	0.02

^a n = 188

^b - Sample size for average catches per successful trapper

Table 25. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 2000-01 season.

Species	Harvest ^a	SE	Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper ^b	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	284	84.96	0.97	0.29	6.24	1.73	25	15.53	0.02
Raccoon	9,809	899.19	33.26	3.02	36.91	3.27	146	90.68	0.01
Muskrat	914	534.02	3.12	1.82	21.83	12.67	23	14.29	0.02
Red Fox	173	32.15	0.59	0.11	3.06	0.47	31	19.26	0.02
Gray Fox	544	55.97	1.96	0.19	4.60	0.36	65	40.37	0.03
Bobcat	950	81.68	3.19	0.27	5.49	0.40	95	59.01	0.03
Opossum	3,524	330.78	11.77	1.11	18.44	1.55	105	65.22	0.03
Otter	1,418	106.75	4.76	0.36	8.20	0.48	95	59.01	0.03
Spotted Skunk	62	14.98	0.21	0.05	3.78	0.41	9	5.59	0.01
Striped Skunk	679	101.71	2.28	0.34	6.54	0.86	57	35.40	0.03
Coyote	2,818	456.79	9.42	1.55	17.20	2.66	90	55.90	0.03
Weasel	2	1.22	0.01	0	1.00	N/A	1	0.62	0.00
Nutria	1,345	610.47	4.58	2.08	16.07	7.22	46	28.57	0.02
Beaver	8,730	923.53	29.52	3.12	39.31	3.94	122	75.78	0.02

^a n = 178

^b n = 161

^c Sample size for average catches per successful trapper

Table 26. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trapper, and percent successful trappers by furbearer species for the 2001-02 season.

Species	Harvest ^a	SE	Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper ^b	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	65	19.91	0.22	0.07	2.38	0.59	13	9.22	0.02
Raccoon	9,172	687.85	30.12	2.28	36.26	2.48	121	85.82	0.02
Muskrat	307	72.22	1.04	0.24	4.90	1.00	30	21.28	0.02
Red Fox	111	28.45	0.38	0.10	2.65	0.56	20	14.18	0.02
Gray Fox	725	108.67	2.43	0.37	6.43	0.82	54	38.30	0.03
Bobcat	1,125	110.31	3.74	0.37	6.56	0.53	82	58.16	0.03
Opossum	3,516	416.54	11.80	1.40	18.48	1.99	91	64.54	0.03
Otter	1,651	151.47	5.44	0.50	9.63	0.71	82	58.16	0.03
Spotted Skunk	42	15.84	0.14	0.05	3.33	0.89	6	4.26	0.01
Striped Skunk	761	116.23	2.55	0.39	6.74	0.88	54	38.30	0.03
Coyote	2,659	331.61	8.69	1.10	14.96	1.70	85	60.24	0.03
Weasel	0	0	0	0	N/A	0	N/A	N/A	N/A
Nutria	1,658	763.26	2.06	0.37	20.87	9.46	38	26.95	0.03
Beaver	8,495	837.75	28.14	2.80	36.61	3.35	111	78.72	0.02

^a n = 155

^b n = 141

^c Sample size for average catches per successful trapper

Table 27. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 2002-03 season.

Species	Harvest ^a	S E	Catch Per Trapper ^a	S E	Average Catch Per Successful Trapper ^b	S E	n ^c	Percent Successful Trappers ^b	S E
Mink	408	102.29	1.25	0.32	6.29	1.50	42	20.59	0.02
Raccoon	12,185	837.64	35.96	2.58	41.04	2.70	192	94.12	0.01
Muskrat	1,117	444.33	3.47	1.41	13.88	5.47	52	25.49	0.02
Red Fox	320	57.93	0.95	0.18	3.98	0.66	52	25.49	0.02
Gray Fox	1,022	81.10	2.99	0.44	7.18	0.45	92	45.10	0.02
Bobcat	1,370	101.24	4.28	0.31	6.87	0.44	129	63.24	0.02
Opossum	6,451	452.11	18.18	1.29	28.57	1.79	146	71.57	0.02
Otter	2,713	207.51	8.42	0.64	11.85	0.83	148	72.55	0.02
Spotted Skunk	339	162.32	1.02	0.51	16.85	7.90	13	6.37	0.01
Striped Skunk	1,225	160.79	2.77	0.25	9.00	1.09	88	43.14	0.02
Coyote	2,818	279.36	8.04	0.85	14.69	1.35	124	60.78	0.02
Weasel	8	4.60	0.02	0.01	5.00	N/A	1	0.49	0.03
Nutria	1,033	172.31	3.01	0.54	8.46	1.34	79	38.73	0.02
Beaver	12,626	853.49	37.53	2.62	44.13	2.82	185	90.69	0.01

^a n =382

^b n =271

^c Sample size for average catches per successful trapper