

Table 7. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trapper, and percent successful trappers by furbearer species for the 2014-15 season. There were 919 trappers who purchased a license for the 2014-15 season. Correcting for the percentage of license holders who indicated they did not trap, and those who trapped feral hogs only, there was an estimated N=711 licensed furbearer trappers in Mississippi in 2014-15 to which estimates were extrapolated.

Species	Harvest ^a	SE	Average Catch Per Trapper ^a	SE	Average Catch Per Successful Trapper	SE	N ^b	Percent Successful Trappers ^a	SE
Mink	95	19.42	0.13	0.02	3.00	0.51	22	4.49	0.52
Raccoon	13,804	507.67	19.42	0.71	23.44	0.82	405	82.82	0.95
Muskrat	508	54.11	0.72	0.08	6.03	0.49	58	11.86	0.82
Red Fox	350	27.61	0.49	0.04	2.32	0.14	104	21.27	1.04
Gray Fox	1,860	97.74	2.62	0.14	5.90	0.26	217	44.38	1.26
Bobcat	2,175	117.50	3.06	0.17	6.68	0.31	224	45.81	1.26
Opossum	7,206	352.21	10.14	0.50	16.91	0.75	293	59.92	1.24
Otter	1,745	112.44	2.45	0.16	6.22	0.35	193	39.47	1.24
Spotted Skunk	44	10.10	0.06	0.01	3.00	0.48	10	2.05	0.36
Striped Skunk	654	49.74	0.92	0.07	4.10	0.25	110	22.50	1.06
Coyote	5,173	409.68	7.27	0.58	13.68	1.03	260	53.17	1.26
Weasel	1	0.81	0.00	0.00	1.00	0.00	1	0.20	0.11
Nutria	842	111.84	1.18	0.16	6.09	0.75	95	19.43	0.10
Beaver	8,453	394.74	11.89	0.56	18.58	0.79	313	64.00	1.21

^a n = 489

^b n=sample size for average catches per successful trapper