

ANNUAL HARVEST SUMMARY REPORT

DEER MANAGEMENT ASSISTANCE PROGRAM

Chickasawhay WMA Jones Co

2024-2025 Season



The goal for this club/cooperator based on the most recent Annual DMAP Application completed is as follows:

Maximize Recreational Harvest

A description of this goal, as well as other deer herd management goals may be found in Section 8 of the Annual DMAP Application. To view your Annual DMAP Application contact your DMAP Biologist or visit <https://xnet2.mdwfp.com/NRIS/>.

DMAP VISION

To put the landowner/cooperator in a better position to manage their lands for a healthy deer herd, while maintaining habitat integrity.



DMAP REPORT

For more information, visit our website
www.mdwfp.com/dmap



DMAP Harvest Summary

Date: 09/02/2025

*** Chickasawhay WMA Jones Co ***

Club ID: 7007

Season: 2024	Acres: 30000
Location: Jones County, MS	Years on DMAP: 34
Soil Region: Lower Coastal Plain	Biologist: Blake Stefano
	% Data Collected: 100

Table 1. Harvest Summary

	Number Harvested	Acres Per Deer Harvested
Bucks	37	811
Does	16	1875
Totals	53	566

Table 2. Harvest History

	2024	2023	2022	2021	2020	2019	2018	2017	2016	2015	RG 2023	Club Avg
Acres	30000	30000	30000	30000	N/A	30000	30000	30000	30000	30000	237285	30000
Total Deer	53	10	27	19	N/A	28	22	33	39	60	922	32
Buck	37	7	17	13	N/A	18	19	27	17	30	457	21
Doe	16	3	10	6	N/A	10	3	6	22	30	465	12
Acres/Deer	566	3000	1111	1579	N/A	1071	1364	909	769	500	257	928
Acres/Buck	811	4286	1765	2308	N/A	1667	1579	1111	1765	1000	519	1459
Acres/Doe	1875	10000	3000	5000	N/A	3000	10000	5000	1364	1000	510	2547

Table 3. Age Distribution

	Bucks		Does		Management Bucks		Criteria Bucks	
Age	#	%	#	%	#	%	#	%
0.5	0	0	3	19	0	0	0	0
1.5	1	3	1	6	0	0	1	3
2.5	1	3	0	0	0	0	1	3
3.5	8	22	5	31	0	0	8	22
4.5	15	41	2	12	0	0	15	41
5.5	6	16	1	6	0	0	6	16
6.5	4	11	4	25	0	0	4	11
7.5	2	5	0	0	0	0	2	5
8.5	0	0	0	0	0	0	0	0

DMAP Harvest Summary

Date: 09/02/2025

*** Chickasawhay WMA Jones Co ***

Club ID: 7007

Table 4. Statistics for Doe Harvest

Age	Number Harvested	Avg Live Weight	Percent Lactating
0.5	3	55	0
1.5	1	80	100
2.5	0	0	0
3.5	5	88	60
4.5	2	92	100
5.5	1	84	100
6.5	4	86	0
7.5	0	0	0
8.5	0	0	0
Totals	16	81	

Table 5. Doe Harvest History

	2024	2023	2022	2021	2020	2019	2018	2017	2016	2015	RG 2023	Club Avg
Total Does	16	2	10	6	8	10	3	6	21	27	443	11
Acres/Doe	1875	15000	3000	5000	3750	3000	10000	5000	1429	1111	583	2752
% Harvest By Age												
% 0.5	19	50	30	50	0	0	0	0	5	7	5	12
% 1.5	6	0	30	0	0	10	0	0	24	11	14	12
% 2.5	0	0	10	0	38	10	0	50	14	7	11	12
% 3.5+	75	50	30	50	62	80	100	50	57	74	68	64
Average Weight												
0.5	55	57	81	65	0	0	0	0	40	38	57	60
1.5	80	0	73	0	0	55	0	0	67	73	79	70
2.5	0	0	76	0	73	80	0	70	72	86	92	75
3.5+	88	89	96	103	78	84	94	87	82	86	94	85
% Lactation												
% 1.5	100	0	0	0	0	0	0	0	0	0	3	8
% 2.5	0	0	0	0	33	0	0	33	0	50	30	23
% 2.5+	50	0	50	33	38	33	67	33	7	27	49	31
% 3.5+	50	0	67	33	40	38	67	33	8	25	52	33

DMAP Harvest Summary

Date: 09/02/2025

*** Chickasawhay WMA Jones Co ***

Club ID: 7007

Table 6. Statistics for Yearling Bucks

	Number	Percent	Avg. Live Weight	Average Points
Spikes	0	0	0	0
Forks	0	0	0	0
Totals	1			

Table 7. Buck Statistics

	Live Weight		Points		Base		Length		Spread	
Age	Number	Average	Number	Average	Number	Average	Number	Average	Number	Average
0.5	0	0	0	N/A	0	N/A	0	N/A	0	N/A
1.5	1	74	1	0	0	N/A	0	N/A	0	N/A
2.5	1	150	1	6	1	2.6	1	15.4	1	15.0
3.5	8	124	8	5.6	8	2.9	8	12.5	8	10.7
4.5	15	131	15	6.5	15	3.3	15	14.7	15	11.9
5.5	6	139	6	5.8	6	3.0	6	15.2	6	12.6
6.5	4	136	4	7	4	3.7	4	14.5	4	11.9
7.5	2	142	2	7	2	4.0	2	14.9	2	12.4
8.5	0	0	0	N/A	0	N/A	0	N/A	0	N/A

Date: 09/02/2025

*** Chickasawhay WMA Jones Co ***

Club ID: 7007

Table 8. Buck Harvest History (All Bucks)

	2024	2023	2022	2021	2020	2019	2018	2017	2016	2015	RG 2023	Club Avg
Bucks	37	7	17	13	27	18	19	27	15	30	445	21
Acres/Buck	811	4286	1765	2308	1111	1667	1579	1111	2000	1000	581	1429
% .5 Bucks	0	0	0	0	0	0	0	0	0	0	2	0
# .5 Bucks	0	0	0	0	0	0	0	0	0	0	8	0
Avg Weight	0	0	0	0	0	0	0	0	0	0	63	0
% 1.5 Bucks	3	0	0	8	0	0	0	0	7	10	8	3
# 1.5 Bucks	1	0	0	1	0	0	0	0	1	3	35	0.6
% Spikes	0	0	0	100	0	0	0	0	0	67	51	50
Avg Weight	74	0	0	115	0	0	0	0	63	95	95	90
Avg Points	0	0	0	2	0	0	0	0	0	1.33	2	1
Avg Base	0	0	0	0	0	0	0	0	0	1.8	2.1	1.8
Avg Length	0	0	0	0	0	0	0	0	0	7.5	6.4	7.5
Avg Spread	0	0	0	0	0	0	0	0	0	6.0	6.4	6.0
% 2.5 Bucks	3	0	0	8	0	17	21	37	13	3	9	10
# 2.5 Bucks	1	0	0	1	0	3	4	10	2	1	39	2.2
Avg Weight	150	0	0	139	0	105	138	117	114	130	126	122
Avg Points	6	0	0	4	0	3.33	5.5	5.8	7.5	8	6	6
Avg Base	2.6	0	0	2.6	0	2.6	2.9	3.4	2.9	3.5	2.8	3.1
Avg Length	15.4	0	0	11.3	0	11.2	13.8	14.6	13.6	13.2	12.9	13.7
Avg Spread	15.0	0	0	10.0	0	10.2	11.2	12.0	11.0	11.5	10.7	11.6
% 3.5 Bucks	22	14	35	54	41	28	63	26	33	20	24	32
# 3.5 Bucks	8	1	6	7	11	5	12	7	5	6	109	6.8
Avg Weight	124	127	143	139	135	147	133	126	119	123	138	132
Avg Points	5.62	5	6.67	6.29	7.09	6.6	6.75	6.29	4.8	6.5	7	6
Avg Base	2.9	3.5	3.3	3.2	3.4	3.2	3.3	2.9	3.2	3.5	3.3	3.2
Avg Length	12.5	15.0	16.2	14.8	15.3	14.9	15.0	13.8	13.0	13.7	14.3	14.4
Avg Spread	10.7	12.0	13.1	11.6	12.9	12.3	11.8	11.4	11.2	10.5	11.5	11.8
% 4.5 + Bucks	73	86	65	31	59	56	16	37	47	67	53	54
# 4.5 + Bucks	27	6	11	4	16	10	3	10	7	20	122	11.4
Avg Weight	135	136	138	146	136	146	138	151	140	131	144	138
Avg Points	6.44	7.67	6.36	7.5	7.06	7.4	8	6.6	7.14	6.6	8	7
Avg Base	3.4	3.1	3.4	3.6	3.6	3.8	4.0	3.8	3.7	3.6	3.7	3.7
Avg Length	14.8	15.8	15.3	16.9	15.9	16.0	18.2	16.5	16.1	14.9	16.3	15.6
Avg Spread	12.1	13.1	12.3	12.9	12.6	13.2	14.8	13.3	13.5	12.5	13.0	12.7

Date: 09/02/2025

*** Chickasawhay WMA Jones Co ***

Club ID: 7007

Table 9. Criteria Buck Harvest History

	2024	2023	2022	2021	2020	2019	2018	2017	2016	2015	RG 2023	Club Avg
Bucks	37	7	17	13	27	18	19	27	15	30	432	21
Acres/Buck	811	4286	1765	2308	1111	1667	1579	1111	2000	1000	598	1429
% .5 Bucks	0	0	0	0	0	0	0	0	0	0	2	0
# .5 Bucks	0	0	0	0	0	0	0	0	0	0	8	0
Avg Weight	0	0	0	0	0	0	0	0	0	0	63	0
% 1.5 Bucks	3	0	0	8	0	0	0	0	7	10	8	3
# 1.5 Bucks	1	0	0	1	0	0	0	0	1	3	33	0.6
% Spikes	0	0	0	100	0	0	0	0	0	67	48	50
Avg Weight	74	0	0	115	0	0	0	0	63	95	95	90
Avg Points	0	0	0	2	0	0	0	0	0	1.33	2	1
Avg Base	0	0	0	0	0	0	0	0	0	1.8	2.1	1.8
Avg Length	0	0	0	0	0	0	0	0	0	7.5	6.6	7.5
Avg Spread	0	0	0	0	0	0	0	0	0	6.0	6.4	6.0
% 2.5 Bucks	3	0	0	8	0	17	21	37	13	3	9	10
# 2.5 Bucks	1	0	0	1	0	3	4	10	2	1	38	2.2
Avg Weight	150	0	0	139	0	105	138	117	114	130	126	122
Avg Points	6	0	0	4	0	3.33	5.5	5.8	7.5	8	6	6
Avg Base	2.6	0	0	2.6	0	2.6	2.9	3.4	2.9	3.5	2.8	3.1
Avg Length	15.4	0	0	11.3	0	11.2	13.8	14.6	13.6	13.2	12.9	13.7
Avg Spread	15.0	0	0	10.0	0	10.2	11.2	12.0	11.0	11.5	10.7	11.6
% 3.5 Bucks	22	14	35	54	41	28	63	26	33	20	24	32
# 3.5 Bucks	8	1	6	7	11	5	12	7	5	6	105	6.8
Avg Weight	124	127	143	139	135	147	133	126	119	123	138	132
Avg Points	5.62	5	6.67	6.29	7.09	6.6	6.75	6.29	4.8	6.5	7	6
Avg Base	2.9	3.5	3.3	3.2	3.4	3.2	3.3	2.9	3.2	3.5	3.3	3.2
Avg Length	12.5	15.0	16.2	14.8	15.3	14.9	15.0	13.8	13.0	13.7	14.3	14.4
Avg Spread	10.7	12.0	13.1	11.6	12.9	12.3	11.8	11.4	11.2	10.5	11.5	11.8
% 4.5 + Bucks	73	86	65	31	59	56	16	37	47	67	54	54
# 4.5 + Bucks	27	6	11	4	16	10	3	10	7	20	118	11.4
Avg Weight	135	136	138	146	136	146	138	151	140	131	143	138
Avg Points	6.44	7.67	6.36	7.5	7.06	7.4	8	6.6	7.14	6.6	8	7
Avg Base	3.4	3.1	3.4	3.6	3.6	3.8	4.0	3.8	3.7	3.6	3.7	3.7
Avg Length	14.8	15.8	15.3	16.9	15.9	16.0	18.2	16.5	16.1	14.9	16.3	15.6
Avg Spread	12.1	13.1	12.3	12.9	12.6	13.2	14.8	13.3	13.5	12.5	13.0	12.7

DMAP Harvest Summary

Date: 09/02/2025

*** Chickasawhay WMA Jones Co ***

Club ID: 7007

Table 10. Management Buck Harvest History

[illegible]

Description of the Data and Glossary of Terms

This report is based on data MDWFP receives from DMAP clubs. The club/cooperator information in the heading, including acres under management and soil region, is based on the Annual DMAP Application. Please contact your DMAP Biologist if any of the information in the heading is incorrect. The “% Data Collected” field is the percentage of deer reported on the DMAP datasheet for which jawbones were received. Harvest totals in Tables 1 and 2 are based on deer harvest reported on the DMAP datasheet (regardless of whether a jawbone was received). All other tables are based on submitted jawbones in adequate condition to age using the tooth wear and replacement technique. For all tables including Soil Region Average data, the regional data column is labeled “RG20xx.” For all tables, Criteria Buck and Management Buck are defined below. Buck designations are made by the DMAP Biologist. If no designations are made, all bucks are classified as Criteria. If you have questions about Management and Criteria bucks, please contact your DMAP Biologist.

Soil Region Average (e.g., Regional Average or RG20xx) – the average of all deer in a specified parameter (i.e., harvest total, lactation, weight, etc.) reported for the year in question is used as an indicator of overall deer herd health by comparing club averages to other DMAP clubs in the same soil region.

Criteria Buck or perhaps a “trophy buck” – a buck deemed to have antler characteristics that satisfy the club’s goals for antler quality. This determination is generally made by the DMAP biologist in collaboration with the DMAP club and considers buck age, antler characteristics, soil region averages, and the goals of the DMAP club.

Management Buck or perhaps a “cull buck” – a harvested buck deemed to have below average antler characteristics compared to other bucks in the same age class and soil region. As with Criteria bucks, this determination is generally made by the DMAP biologist in collaboration with the DMAP club.

Lactation Index – the percentage of does for which a DMAP club recorded “detecting lactation” on the datasheet. Where sample sizes are sufficient, this index is used to determine the number of adult does that successfully raised fawns and is used to help estimate fawn recruitment. These data may be highly variable based on the timing of doe harvest in the annual cycle, as well as how thoroughly club members check for lactation. Generally, a favorable lactation index would be 60% or greater for adult does. A low lactation index (with adequate sample size) may indicate poor habitat, overpopulation, or a significant predation influence. Doe harvest early in the season (Oct- Dec) is generally better in estimating fawn recruitment vs late season harvest (after does finish lactating). For this report, lactation index is adjusted to account for late-season harvest.

Dressed Weights - average weight to the nearest pound of a dressed animal. A dressed animal should have both the forward (heart and lungs and rear (intestines) viscera removed but not any part of the remaining carcass. Comparing club average body weights from year to year or to the coordinating soil region average can be an indicator of overall herd health. Typically, doe carcass condition is a more reliable indicator of herd health than that of bucks due to the drastic fluctuation of buck body weights during the rut. Significant annual changes in carcass condition may indicate changes in habitat, mast crop, herd density, disease influence, or even severe weather conditions.

Age Distribution of Harvest – an estimate of age structure of the population, though generally only the doe segment is assessed, as buck harvest is often biased. Balanced deer herds typically show some harvest across all age classes. However, an age structure with more older animals (>40% 3 years old or greater) may indicate a stockpile of older animals which could signal an overabundant deer herd. Age distribution data is generally best considered in context with other harvest data (i.e., lactation index, doe body weights), as well as observation data (i.e., adult sex ratio, fawn recruitment) and even camera survey data (i.e., deer density, buck age structure).

Fawn Recruitment – a measurement of the number of fawns per adult doe that survive to at least 4-6 months of age. Recruitment tends to improve with increased harvest due to freeing up available resources. Generally fawn recruitment is measured using hunter observation data or camera surveys.

Adult Sex Ratio – ratio of the total number of does to total number of bucks in a deer herd. Typically, adult sex ratio is measured using observation data or camera surveys. Deer are born at nearly a 1:1 ratio of does to bucks. Thus, adult sex ratios closer to 1:1 are ideal. In trophy management scenarios adult sex ratio may be skewed towards more bucks, but most deer herds range from 1.5:1 to 3:1.

Important DMAP Reminders for Each Season

- For more information, visit www.mdwfp.com/DMAP and check out the Deer Managers Toolkit.
- If club acreages are incorrect, they should be updated on your DMAP Application. Contact your DMAP Biologist for help updating your DMAP Application.
- Calibrate your body weight scale(s) each year, especially if your club has multiple scales.
- Don't forget to retrieve jawbones from taxidermy bucks. The mature bucks are the most important to have aged.
- Do your best to collect all data completely. Check udders thoroughly for lactation.
- Hunter observation data can provide useful information, especially in clubs with small harvest data sets. Ask your biologist about hunter observation books.