

ANNUAL HARVEST SUMMARY REPORT

DEER MANAGEMENT ASSISTANCE PROGRAM

Marion County WMA

2024-2025 Season



The goal for this club/cooperator based on the most recent Annual DMAP Application completed is as follows:

A description of this goal, as well as other deer herd management goals may be found in Section 8 of the Annual DMAP Application. To view your Annual DMAP Application contact your DMAP Biologist or visit <https://xnet2.mdwfp.com/NRIS/>.

DMAP VISION

To put the landowner/cooperator in a better position to manage their lands for a healthy deer herd, while maintaining habitat integrity.



DMAP REPORT

For more information, visit our website
www.mdwfp.com/dmap



DMAP Harvest Summary

Date: 08/27/2025

*** Marion County WMA ***

Club ID: 7013

Season: 2024	Acres: 7125
Location: Marion County, MS	Years on DMAP: 32
Soil Region: Lower Coastal Plain	Biologist: Blake Stefano
	% Data Collected: 96

Table 1. Harvest Summary

	Number Harvested	Acres Per Deer Harvested
Bucks	28	254
Does	18	396
Totals	46	155

Table 2. Harvest History

	2024	2023	2022	2021	2020	2019	2018	2017	2016	2015	RG 2023	Club Avg
Acres	7125	7125	7125	7125	7125	7125	7125	7125	7125	7125	237285	7125
Total Deer	46	55	48	52	58	63	48	49	50	51	922	52
Buck	28	28	19	20	23	34	20	19	22	18	457	23
Doe	18	27	29	32	35	29	28	30	28	33	465	29
Acres/Deer	155	130	148	137	123	113	148	145	142	140	257	137
Acres/Buck	254	254	375	356	310	210	356	375	324	396	519	308
Acres/Doe	396	264	246	223	204	246	254	238	254	216	510	247

Table 3. Age Distribution

	Bucks		Does		Management Bucks		Criteria Bucks	
Age	#	%	#	%	#	%	#	%
0.5	0	0	0	0	0	0	0	0
1.5	0	0	1	6	0	0	0	0
2.5	6	22	0	0	0	0	6	22
3.5	9	33	5	29	0	0	9	33
4.5	6	22	4	24	0	0	6	22
5.5	3	11	1	6	0	0	3	11
6.5	2	7	3	18	0	0	2	7
7.5	1	4	2	12	0	0	1	4
8.5	0	0	1	6	0	0	0	0

DMAP Harvest Summary

Date: 08/27/2025

*** Marion County WMA ***

Club ID: 7013

Table 4. Statistics for Doe Harvest

Age	Number Harvested	Avg Live Weight	Percent Lactating
0.5	0	0	0
1.5	1	50	0
2.5	0	0	0
3.5	5	83	40
4.5	4	82	25
5.5	1	80	0
6.5	3	95	100
7.5	2	110	0
8.5	1	122	0
Totals	17	89	

Table 5. Doe Harvest History

	2024	2023	2022	2021	2020	2019	2018	2017	2016	2015	RG 2023	Club Avg
Total Does	17	26	24	31	32	29	26	30	26	33	443	27
Acres/Doe	419	274	297	230	223	246	274	238	274	216	583	260
% Harvest By Age												
% 0.5	0	23	4	6	6	3	0	3	15	15	5	8
% 1.5	6	19	21	26	12	21	38	30	15	12	14	20
% 2.5	0	12	21	16	16	28	19	13	12	9	11	15
% 3.5+	94	46	54	52	66	48	42	53	58	64	68	57
Average Weight												
0.5	0	39	60	50	35	50	0	60	50	52	57	47
1.5	50	62	80	68	68	69	72	70	79	72	79	70
2.5	0	74	73	78	70	81	84	78	80	90	92	78
3.5+	90	81	91	88	83	87	90	88	84	94	94	84
% Lactation												
% 1.5	0	0	0	0	0	0	0	11	0	0	3	2
% 2.5	0	33	40	60	60	50	40	75	100	33	30	54
% 2.5+	38	40	56	52	80	55	56	60	78	79	49	61
% 3.5+	38	42	62	50	85	57	64	56	73	86	52	63

DMAP Harvest Summary

Date: 08/27/2025

*** Marion County WMA ***

Club ID: 7013

Table 6. Statistics for Yearling Bucks

	Number	Percent	Avg. Live Weight	Average Points
Spikes	0	0	0	0
Forks	0	0	0	0
Totals	0			

Table 7. Buck Statistics

	Live Weight		Points		Base		Length		Spread	
Age	Number	Average	Number	Average	Number	Average	Number	Average	Number	Average
0.5	0	0	0	N/A	0	N/A	0	N/A	0	N/A
1.5	0	0	0	N/A	0	N/A	0	N/A	0	N/A
2.5	6	100	6	3	6	2.0	5	8.6	2	11.9
3.5	9	134	9	6.1	8	3.5	8	13.3	7	12.0
4.5	6	134	6	7.5	6	3.3	6	15.1	6	11.9
5.5	3	150	3	8.3	3	4.0	3	18.1	3	13.4
6.5	2	114	2	5	2	2.8	2	9.1	1	12.2
7.5	1	170	1	9	1	3.5	1	16.9	1	12.0
8.5	0	0	0	N/A	0	N/A	0	N/A	0	N/A

DMAP Harvest Summary

Date: 08/27/2025

*** Marion County WMA ***

Club ID: 7013

Table 8. Buck Harvest History (All Bucks)

	2024	2023	2022	2021	2020	2019	2018	2017	2016	2015	RG 2023	Club Avg
Bucks	27	24	18	20	21	30	16	19	21	17	445	21
Acres/Buck	264	297	396	356	339	238	445	375	339	419	581	335
% .5 Bucks	0	0	0	5	10	3	12	5	10	6	2	5
# .5 Bucks	0	0	0	1	2	1	2	1	2	1	8	10
Avg Weight	0	0	0	40	42	80	47	50	59	0	63	52
% 1.5 Bucks	0	25	22	25	29	20	19	11	19	18	8	18
# 1.5 Bucks	0	6	4	5	6	6	3	2	4	3	35	3.9
% Spikes	0	33	50	100	50	67	67	0	75	67	51	59
Avg Weight	0	72	88	93	82	76	71	89	92	93	95	83
Avg Points	0	2.33	2	2	2	1.5	2	2	2.25	2.33	2	1
Avg Base	0	2.1	0.6	1.5	2.1	1.2	1.4	2.6	1.7	2.0	2.1	1.7
Avg Length	0	2.5	1.5	3.5	2.5	2.7	0.9	11.2	4.8	4.2	6.4	3.3
Avg Spread	0	3.2	4.0	4.9	3.9	3.5	3.2	8.5	5.7	4.6	6.4	4.6
% 2.5 Bucks	22	17	50	20	14	30	38	16	14	6	9	23
# 2.5 Bucks	6	4	9	4	3	9	6	3	3	1	39	4.8
Avg Weight	100	112	121	125	120	119	126	127	130	118	126	119
Avg Points	3	6	5.33	6.5	5.33	6.67	5.67	6	5	3	6	6
Avg Base	2.2	3.1	3.3	3.0	2.8	3.4	3.1	3.2	3.2	2.8	2.8	3.1
Avg Length	8.6	13.6	11.6	13.0	13.5	14.3	13.7	14.1	11.0	12.4	12.9	12.6
Avg Spread	11.9	10.9	10.3	11.6	13.1	11.1	11.2	11.0	12.0	8.5	10.7	11.1
% 3.5 Bucks	33	33	17	15	33	20	25	53	19	47	24	29
# 3.5 Bucks	9	8	3	3	7	6	4	10	4	8	109	6.2
Avg Weight	134	125	127	123	139	133	137	132	126	128	138	131
Avg Points	6.11	6	6.67	6.67	7.43	7.17	5.5	5.9	7.25	7.25	7	7
Avg Base	3.8	3.2	3.4	3.3	3.6	4.0	3.4	3.6	3.5	3.3	3.3	3.5
Avg Length	13.3	13.3	15.2	14.8	16.6	16.1	14.5	15.3	16.2	15.7	14.3	15.0
Avg Spread	12.0	11.3	12.8	12.2	12.9	12.8	10.1	11.5	12.3	13.5	11.5	12.2
% 4.5 + Bucks	44	25	11	35	14	27	6	16	38	24	53	25
# 4.5 + Bucks	12	6	2	7	3	8	1	3	8	4	122	5.4
Avg Weight	137	130	118	133	130	150	125	135	153	132	144	138
Avg Points	7.42	7.83	7	7.14	6.33	7.5	6	9	7.25	8.25	8	8
Avg Base	3.4	3.8	3.8	3.7	3.9	4.0	3.8	3.8	4.4	3.8	3.7	3.7
Avg Length	15.0	15.3	17.0	15.2	16.7	17.3	16.2	18.5	17.0	17.4	16.3	16.3
Avg Spread	12.4	11.9	13.0	11.7	13.7	14.0	12.0	15.2	13.8	14.5	13.0	13.1

DMAP Harvest Summary

Date: 08/27/2025

*** Marion County WMA ***

Club ID: 7013

Table 9. Criteria Buck Harvest History

	2024	2023	2022	2021	2020	2019	2018	2017	2016	2015	RG 2023	Club Avg
Bucks	27	24	18	20	21	30	16	19	21	17	432	21
Acres/Buck	264	297	396	356	339	238	445	375	339	419	598	335
% .5 Bucks	0	0	0	5	10	3	12	5	10	6	2	5
# .5 Bucks	0	0	0	1	2	1	2	1	2	1	8	10
Avg Weight	0	0	0	40	42	80	47	50	59	0	63	52
% 1.5 Bucks	0	25	22	25	29	20	19	11	19	18	8	18
# 1.5 Bucks	0	6	4	5	6	6	3	2	4	3	33	3.9
% Spikes	0	33	50	100	50	67	67	0	75	67	48	59
Avg Weight	0	72	88	93	82	76	71	89	92	93	95	83
Avg Points	0	2.33	2	2	2	1.5	2	2	2.25	2.33	2	1
Avg Base	0	2.1	0.6	1.5	2.1	1.2	1.4	2.6	1.7	2.0	2.1	1.7
Avg Length	0	2.5	1.5	3.5	2.5	2.7	0.9	11.2	4.8	4.2	6.6	3.3
Avg Spread	0	3.2	4.0	4.9	3.9	3.5	3.2	8.5	5.7	4.6	6.4	4.6
% 2.5 Bucks	22	17	50	20	14	30	38	16	14	6	9	23
# 2.5 Bucks	6	4	9	4	3	9	6	3	3	1	38	4.8
Avg Weight	100	112	121	125	120	119	126	127	130	118	126	119
Avg Points	3	6	5.33	6.5	5.33	6.67	5.67	6	5	3	6	6
Avg Base	2.2	3.1	3.3	3.0	2.8	3.4	3.1	3.2	3.2	2.8	2.8	3.1
Avg Length	8.6	13.6	11.6	13.0	13.5	14.3	13.7	14.1	11.0	12.4	12.9	12.6
Avg Spread	11.9	10.9	10.3	11.6	13.1	11.1	11.2	11.0	12.0	8.5	10.7	11.1
% 3.5 Bucks	33	33	17	15	33	20	25	53	19	47	24	29
# 3.5 Bucks	9	8	3	3	7	6	4	10	4	8	105	6.2
Avg Weight	134	125	127	123	139	133	137	132	126	128	138	131
Avg Points	6.11	6	6.67	6.67	7.43	7.17	5.5	5.9	7.25	7.25	7	7
Avg Base	3.8	3.2	3.4	3.3	3.6	4.0	3.4	3.6	3.5	3.3	3.3	3.5
Avg Length	13.3	13.3	15.2	14.8	16.6	16.1	14.5	15.3	16.2	15.7	14.3	15.0
Avg Spread	12.0	11.3	12.8	12.2	12.9	12.8	10.1	11.5	12.3	13.5	11.5	12.2
% 4.5 + Bucks	44	25	11	35	14	27	6	16	38	24	54	25
# 4.5 + Bucks	12	6	2	7	3	8	1	3	8	4	118	5.4
Avg Weight	137	130	118	133	130	150	125	135	153	132	143	138
Avg Points	7.42	7.83	7	7.14	6.33	7.5	6	9	7.25	8.25	8	8
Avg Base	3.4	3.8	3.8	3.7	3.9	4.0	3.8	3.8	4.4	3.8	3.7	3.7
Avg Length	15.0	15.3	17.0	15.2	16.7	17.3	16.2	18.5	17.0	17.4	16.3	16.3
Avg Spread	12.4	11.9	13.0	11.7	13.7	14.0	12.0	15.2	13.8	14.5	13.0	13.1

DMAP Harvest Summary

Date: 08/27/2025

*** Marion County WMA ***

Club ID: 7013

Table 10. Management Buck Harvest History

[illegible]

Description of the Data and Glossary of Terms

This report is based on data MDWFP receives from DMAP clubs. The club/cooperator information in the heading, including acres under management and soil region, is based on the Annual DMAP Application. Please contact your DMAP Biologist if any of the information in the heading is incorrect. The “% Data Collected” field is the percentage of deer reported on the DMAP datasheet for which jawbones were received. Harvest totals in Tables 1 and 2 are based on deer harvest reported on the DMAP datasheet (regardless of whether a jawbone was received). All other tables are based on submitted jawbones in adequate condition to age using the tooth wear and replacement technique. For all tables including Soil Region Average data, the regional data column is labeled “RG20xx.” For all tables, Criteria Buck and Management Buck are defined below. Buck designations are made by the DMAP Biologist. If no designations are made, all bucks are classified as Criteria. If you have questions about Management and Criteria bucks, please contact your DMAP Biologist.

Soil Region Average (e.g., Regional Average or RG20xx) – the average of all deer in a specified parameter (i.e., harvest total, lactation, weight, etc.) reported for the year in question is used as an indicator of overall deer herd health by comparing club averages to other DMAP clubs in the same soil region.

Criteria Buck or perhaps a “trophy buck” – a buck deemed to have antler characteristics that satisfy the club’s goals for antler quality. This determination is generally made by the DMAP biologist in collaboration with the DMAP club and considers buck age, antler characteristics, soil region averages, and the goals of the DMAP club.

Management Buck or perhaps a “cull buck” – a harvested buck deemed to have below average antler characteristics compared to other bucks in the same age class and soil region. As with Criteria bucks, this determination is generally made by the DMAP biologist in collaboration with the DMAP club.

Lactation Index – the percentage of does for which a DMAP club recorded “detecting lactation” on the datasheet. Where sample sizes are sufficient, this index is used to determine the number of adult does that successfully raised fawns and is used to help estimate fawn recruitment. These data may be highly variable based on the timing of doe harvest in the annual cycle, as well as how thoroughly club members check for lactation. Generally, a favorable lactation index would be 60% or greater for adult does. A low lactation index (with adequate sample size) may indicate poor habitat, overpopulation, or a significant predation influence. Doe harvest early in the season (Oct- Dec) is generally better in estimating fawn recruitment vs late season harvest (after does finish lactating). For this report, lactation index is adjusted to account for late-season harvest.

Dressed Weights - average weight to the nearest pound of a dressed animal. A dressed animal should have both the forward (heart and lungs and rear (intestines) viscera removed but not any part of the remaining carcass. Comparing club average body weights from year to year or to the coordinating soil region average can be an indicator of overall herd health. Typically, doe carcass condition is a more reliable indicator of herd health than that of bucks due to the drastic fluctuation of buck body weights during the rut. Significant annual changes in carcass condition may indicate changes in habitat, mast crop, herd density, disease influence, or even severe weather conditions.

Age Distribution of Harvest – an estimate of age structure of the population, though generally only the doe segment is assessed, as buck harvest is often biased. Balanced deer herds typically show some harvest across all age classes. However, an age structure with more older animals (>40% 3 years old or greater) may indicate a stockpile of older animals which could signal an overabundant deer herd. Age distribution data is generally best considered in context with other harvest data (i.e., lactation index, doe body weights), as well as observation data (i.e., adult sex ratio, fawn recruitment) and even camera survey data (i.e., deer density, buck age structure).

Fawn Recruitment – a measurement of the number of fawns per adult doe that survive to at least 4-6 months of age. Recruitment tends to improve with increased harvest due to freeing up available resources. Generally fawn recruitment is measured using hunter observation data or camera surveys.

Adult Sex Ratio – ratio of the total number of does to total number of bucks in a deer herd. Typically, adult sex ratio is measured using observation data or camera surveys. Deer are born at nearly a 1:1 ratio of does to bucks. Thus, adult sex ratios closer to 1:1 are ideal. In trophy management scenarios adult sex ratio may be skewed towards more bucks, but most deer herds range from 1.5:1 to 3:1.

Important DMAP Reminders for Each Season

- For more information, visit www.mdwfp.com/DMAP and check out the Deer Managers Toolkit.
- If club acreages are incorrect, they should be updated on your DMAP Application. Contact your DMAP Biologist for help updating your DMAP Application.
- Calibrate your body weight scale(s) each year, especially if your club has multiple scales.
- Don't forget to retrieve jawbones from taxidermy bucks. The mature bucks are the most important to have aged.
- Do your best to collect all data completely. Check udders thoroughly for lactation.
- Hunter observation data can provide useful information, especially in clubs with small harvest data sets. Ask your biologist about hunter observation books.