

ANNUAL HARVEST SUMMARY REPORT

DEER MANAGEMENT ASSISTANCE PROGRAM

Wolf River WMA

2024-2025 Season



The goal for this club/cooperator based on the most recent Annual DMAP Application completed is as follows:

A description of this goal, as well as other deer herd management goals may be found in Section 8 of the Annual DMAP Application. To view your Annual DMAP Application contact your DMAP Biologist or visit <https://xnet2.mdwfp.com/NRIS/>.

DMAP VISION

To put the landowner/cooperator in a better position to manage their lands for a healthy deer herd, while maintaining habitat integrity.



DMAP REPORT

For more information, visit our website
www.mdwfp.com/dmap



DMAP Harvest Summary

Date: 08/27/2025

*** Wolf River WMA ***

Club ID: 7018

Season: 2024	Acres: 10000
Location: Pearl River County, MS	Years on DMAP: 34
Soil Region: Lower Coastal Plain	Biologist: Blake Stefano
	% Data Collected: 97

Table 1. Harvest Summary

	Number Harvested	Acres Per Deer Harvested
Bucks	23	435
Does	7	1429
Totals	30	333

Table 2. Harvest History

	2024	2023	2022	2021	2020	2019	2018	2017	2016	2015	RG 2023	Club Avg
Acres	10000	10000	10000	10000	10000	10000	10000	10000	10000	10000	237285	10000
Total Deer	30	34	34	28	40	34	41	20	47	49	922	36
Buck	23	19	25	18	19	19	31	19	34	25	457	23
Doe	7	15	9	10	21	15	10	1	13	24	465	12
Acres/Deer	333	294	294	357	250	294	244	500	213	204	257	280
Acres/Buck	435	526	400	556	526	526	323	526	294	400	519	431
Acres/Doe	1429	667	1111	1000	476	667	1000	10000	769	417	510	800

Table 3. Age Distribution

	Bucks		Does		Management Bucks		Criteria Bucks	
Age	#	%	#	%	#	%	#	%
0.5	0	0	0	0	0	0	0	0
1.5	0	0	0	0	0	0	0	0
2.5	4	17	2	33	0	0	4	17
3.5	9	39	1	17	0	0	9	39
4.5	6	26	0	0	0	0	6	26
5.5	3	13	1	17	0	0	3	13
6.5	1	4	1	17	0	0	1	4
7.5	0	0	1	17	0	0	0	0
8.5	0	0	0	0	0	0	0	0

DMAP Harvest Summary

Date: 08/27/2025

*** Wolf River WMA ***

Club ID: 7018

Table 4. Statistics for Doe Harvest

Age	Number Harvested	Avg Live Weight	Percent Lactating
0.5	0	0	0
1.5	0	0	0
2.5	2	90	50
3.5	1	95	100
4.5	0	0	0
5.5	1	105	100
6.5	1	100	0
7.5	1	100	100
8.5	0	0	0
Totals	6	98	

Table 5. Doe Harvest History

	2024	2023	2022	2021	2020	2019	2018	2017	2016	2015	RG 2023	Club Avg
Total Does	6	15	9	10	21	14	9	1	13	24	443	12
Acres/Doe	1667	667	1111	1000	476	714	1111	10000	769	417	583	820
% Harvest By Age												
% 0.5	0	7	0	10	10	0	11	0	8	0	5	5
% 1.5	0	20	11	20	0	29	22	0	23	17	14	16
% 2.5	33	0	0	20	29	14	22	0	0	4	11	12
% 3.5+	67	73	89	50	62	57	44	100	69	79	68	67
Average Weight												
0.5	0	70	0	50	58	0	61	0	60	0	57	59
1.5	0	78	65	82	0	82	83	0	92	88	79	83
2.5	90	0	0	98	88	95	100	0	0	100	92	93
3.5+	100	94	94	100	100	99	99	90	110	102	94	97
% Lactation												
% 1.5	0	0	0	0	0	25	50	0	0	0	3	11
% 2.5	50	0	0	50	33	0	100	0	0	100	30	47
% 2.5+	67	55	25	43	32	50	33	100	67	55	49	47
% 3.5+	75	55	25	40	31	62	0	100	67	53	52	48

DMAP Harvest Summary

Date: 08/27/2025

*** Wolf River WMA ***

Club ID: 7018

Table 6. Statistics for Yearling Bucks

	Number	Percent	Avg. Live Weight	Average Points
Spikes	0	0	0	0
Forks	0	0	0	0
Totals	0			

Table 7. Buck Statistics

	Live Weight		Points		Base		Length		Spread	
Age	Number	Average	Number	Average	Number	Average	Number	Average	Number	Average
0.5	0	0	0	N/A	0	N/A	0	N/A	0	N/A
1.5	0	0	0	N/A	0	N/A	0	N/A	0	N/A
2.5	4	122	4	7.2	4	2.7	4	13.2	4	10.5
3.5	9	130	9	6.6	9	3.4	9	13.8	9	12.6
4.5	6	128	6	7.2	6	3.5	6	15.1	6	12.5
5.5	3	131	3	7.3	3	4.1	3	17.9	3	15.2
6.5	1	160	1	8	1	3.5	1	16.2	1	10.2
7.5	0	0	0	N/A	0	N/A	0	N/A	0	N/A
8.5	0	0	0	N/A	0	N/A	0	N/A	0	N/A

DMAP Harvest Summary

Date: 08/27/2025

*** Wolf River WMA ***

Club ID: 7018

Table 8. Buck Harvest History (All Bucks)

	2024	2023	2022	2021	2020	2019	2018	2017	2016	2015	RG 2023	Club Avg
Bucks	23	18	24	18	19	14	28	17	33	23	445	22
Acres/Buck	435	556	417	556	526	714	357	588	303	435	581	461
% .5 Bucks	0	0	0	0	11	0	0	0	3	0	2	1
# .5 Bucks	0	0	0	0	2	0	0	0	1	0	8	3
Avg Weight	0	0	0	0	51	0	0	0	55	0	63	52
% 1.5 Bucks	0	6	0	0	0	14	0	0	3	4	8	2
# 1.5 Bucks	0	1	0	0	0	2	0	0	1	1	35	0.5
% Spikes	0	100	0	0	0	100	0	0	100	100	51	100
Avg Weight	0	75	0	0	0	100	0	0	100	115	95	98
Avg Points	0	2	0	0	0	2	0	0	2	2	2	2
Avg Base	0	0	0	0	0	2.5	0	0	1.0	0	2.1	1.8
Avg Length	0	0	0	0	0	6.5	0	0	1.0	6.0	6.4	4.5
Avg Spread	0	0	0	0	0	5.0	0	0	0	0	6.4	5.0
% 2.5 Bucks	17	6	29	0	11	7	50	6	27	30	9	21
# 2.5 Bucks	4	1	7	0	2	1	14	1	9	7	39	4.6
Avg Weight	122	110	127	0	115	125	144	115	140	126	126	133
Avg Points	7.25	8	6.14	0	6.5	6	5.71	7	6.22	6.29	6	6
Avg Base	2.7	2.9	3.0	0	2.9	0	3.1	3.5	3.4	3.1	2.8	3.1
Avg Length	13.2	12.8	13.6	0	10.9	8.8	14.1	16.0	13.8	13.2	12.9	13.5
Avg Spread	10.5	10.2	10.8	0	11.3	8.2	11.1	13.0	12.6	11.0	10.7	11.2
% 3.5 Bucks	39	22	33	56	58	64	25	47	42	39	24	41
# 3.5 Bucks	9	4	8	10	11	9	7	8	14	9	109	8.9
Avg Weight	130	130	135	132	148	139	155	149	138	139	138	140
Avg Points	6.56	6.5	6.86	7.1	7.45	6.75	8	7.12	7.21	7.78	7	7
Avg Base	3.4	3.6	3.4	3.5	3.1	3.4	3.7	3.8	3.9	3.6	3.3	3.5
Avg Length	13.8	13.9	14.9	15.1	14.1	14.2	15.9	15.4	15.5	15.2	14.3	14.9
Avg Spread	12.6	10.9	11.7	11.9	11.5	12.6	13.0	13.4	12.0	12.0	11.5	12.2
% 4.5 + Bucks	43	67	38	44	21	14	25	47	24	26	53	34
# 4.5 + Bucks	10	12	9	8	4	2	7	8	8	6	122	7.4
Avg Weight	132	135	138	132	136	175	170	160	160	150	144	146
Avg Points	7.3	7.33	7.56	7.88	7.75	10	7.86	8.12	8.75	8.83	8	8
Avg Base	3.6	3.7	3.7	3.6	4.1	4.2	4.6	4.6	4.4	4.3	3.7	3.7
Avg Length	16.0	16.1	17.1	16.0	16.9	17.4	18.6	18.0	18.7	17.5	16.3	17.1
Avg Spread	13.1	12.5	14.0	12.8	13.4	13.8	14.4	14.8	15.0	14.5	13.0	13.7

DMAP Harvest Summary

Date: 08/27/2025

*** Wolf River WMA ***

Club ID: 7018

Table 9. Criteria Buck Harvest History

	2024	2023	2022	2021	2020	2019	2018	2017	2016	2015	RG 2023	Club Avg
Bucks	23	18	24	18	19	14	28	17	33	23	432	22
Acres/Buck	435	556	417	556	526	714	357	588	303	435	598	461
% .5 Bucks	0	0	0	0	11	0	0	0	3	0	2	1
# .5 Bucks	0	0	0	0	2	0	0	0	1	0	8	3
Avg Weight	0	0	0	0	51	0	0	0	55	0	63	52
% 1.5 Bucks	0	6	0	0	0	14	0	0	3	4	8	2
# 1.5 Bucks	0	1	0	0	0	2	0	0	1	1	33	0.5
% Spikes	0	100	0	0	0	100	0	0	100	100	48	100
Avg Weight	0	75	0	0	0	100	0	0	100	115	95	98
Avg Points	0	2	0	0	0	2	0	0	2	2	2	2
Avg Base	0	0	0	0	0	2.5	0	0	1.0	0	2.1	1.8
Avg Length	0	0	0	0	0	6.5	0	0	1.0	6.0	6.6	4.5
Avg Spread	0	0	0	0	0	5.0	0	0	0	0	6.4	5.0
% 2.5 Bucks	17	6	29	0	11	7	50	6	27	30	9	21
# 2.5 Bucks	4	1	7	0	2	1	14	1	9	7	38	4.6
Avg Weight	122	110	127	0	115	125	144	115	140	126	126	133
Avg Points	7.25	8	6.14	0	6.5	6	5.71	7	6.22	6.29	6	6
Avg Base	2.7	2.9	3.0	0	2.9	0	3.1	3.5	3.4	3.1	2.8	3.1
Avg Length	13.2	12.8	13.6	0	10.9	8.8	14.1	16.0	13.8	13.2	12.9	13.5
Avg Spread	10.5	10.2	10.8	0	11.3	8.2	11.1	13.0	12.6	11.0	10.7	11.2
% 3.5 Bucks	39	22	33	56	58	64	25	47	42	39	24	41
# 3.5 Bucks	9	4	8	10	11	9	7	8	14	9	105	8.9
Avg Weight	130	130	135	132	148	139	155	149	138	139	138	140
Avg Points	6.56	6.5	6.86	7.1	7.45	6.75	8	7.12	7.21	7.78	7	7
Avg Base	3.4	3.6	3.4	3.5	3.1	3.4	3.7	3.8	3.9	3.6	3.3	3.5
Avg Length	13.8	13.9	14.9	15.1	14.1	14.2	15.9	15.4	15.5	15.2	14.3	14.9
Avg Spread	12.6	10.9	11.7	11.9	11.5	12.6	13.0	13.4	12.0	12.0	11.5	12.2
% 4.5 + Bucks	43	67	38	44	21	14	25	47	24	26	54	34
# 4.5 + Bucks	10	12	9	8	4	2	7	8	8	6	118	7.4
Avg Weight	132	135	138	132	136	175	170	160	160	150	143	146
Avg Points	7.3	7.33	7.56	7.88	7.75	10	7.86	8.12	8.75	8.83	8	8
Avg Base	3.6	3.7	3.7	3.6	4.1	4.2	4.6	4.6	4.4	4.3	3.7	3.7
Avg Length	16.0	16.1	17.1	16.0	16.9	17.4	18.6	18.0	18.7	17.5	16.3	17.1
Avg Spread	13.1	12.5	14.0	12.8	13.4	13.8	14.4	14.8	15.0	14.5	13.0	13.7

DMAP Harvest Summary

Date: 08/27/2025

*** Wolf River WMA ***

Club ID: 7018

Table 10. Management Buck Harvest History

[illegible]

Description of the Data and Glossary of Terms

This report is based on data MDWFP receives from DMAP clubs. The club/cooperator information in the heading, including acres under management and soil region, is based on the Annual DMAP Application. Please contact your DMAP Biologist if any of the information in the heading is incorrect. The “% Data Collected” field is the percentage of deer reported on the DMAP datasheet for which jawbones were received. Harvest totals in Tables 1 and 2 are based on deer harvest reported on the DMAP datasheet (regardless of whether a jawbone was received). All other tables are based on submitted jawbones in adequate condition to age using the tooth wear and replacement technique. For all tables including Soil Region Average data, the regional data column is labeled “RG20xx.” For all tables, Criteria Buck and Management Buck are defined below. Buck designations are made by the DMAP Biologist. If no designations are made, all bucks are classified as Criteria. If you have questions about Management and Criteria bucks, please contact your DMAP Biologist.

Soil Region Average (e.g., Regional Average or RG20xx) – the average of all deer in a specified parameter (i.e., harvest total, lactation, weight, etc.) reported for the year in question is used as an indicator of overall deer herd health by comparing club averages to other DMAP clubs in the same soil region.

Criteria Buck or perhaps a “trophy buck” – a buck deemed to have antler characteristics that satisfy the club’s goals for antler quality. This determination is generally made by the DMAP biologist in collaboration with the DMAP club and considers buck age, antler characteristics, soil region averages, and the goals of the DMAP club.

Management Buck or perhaps a “cull buck” – a harvested buck deemed to have below average antler characteristics compared to other bucks in the same age class and soil region. As with Criteria bucks, this determination is generally made by the DMAP biologist in collaboration with the DMAP club.

Lactation Index – the percentage of does for which a DMAP club recorded “detecting lactation” on the datasheet. Where sample sizes are sufficient, this index is used to determine the number of adult does that successfully raised fawns and is used to help estimate fawn recruitment. These data may be highly variable based on the timing of doe harvest in the annual cycle, as well as how thoroughly club members check for lactation. Generally, a favorable lactation index would be 60% or greater for adult does. A low lactation index (with adequate sample size) may indicate poor habitat, overpopulation, or a significant predation influence. Doe harvest early in the season (Oct- Dec) is generally better in estimating fawn recruitment vs late season harvest (after does finish lactating). For this report, lactation index is adjusted to account for late-season harvest.

Dressed Weights - average weight to the nearest pound of a dressed animal. A dressed animal should have both the forward (heart and lungs and rear (intestines) viscera removed but not any part of the remaining carcass. Comparing club average body weights from year to year or to the coordinating soil region average can be an indicator of overall herd health. Typically, doe carcass condition is a more reliable indicator of herd health than that of bucks due to the drastic fluctuation of buck body weights during the rut. Significant annual changes in carcass condition may indicate changes in habitat, mast crop, herd density, disease influence, or even severe weather conditions.

Age Distribution of Harvest – an estimate of age structure of the population, though generally only the doe segment is assessed, as buck harvest is often biased. Balanced deer herds typically show some harvest across all age classes. However, an age structure with more older animals (>40% 3 years old or greater) may indicate a stockpile of older animals which could signal an overabundant deer herd. Age distribution data is generally best considered in context with other harvest data (i.e., lactation index, doe body weights), as well as observation data (i.e., adult sex ratio, fawn recruitment) and even camera survey data (i.e., deer density, buck age structure).

Fawn Recruitment – a measurement of the number of fawns per adult doe that survive to at least 4-6 months of age. Recruitment tends to improve with increased harvest due to freeing up available resources. Generally fawn recruitment is measured using hunter observation data or camera surveys.

Adult Sex Ratio – ratio of the total number of does to total number of bucks in a deer herd. Typically, adult sex ratio is measured using observation data or camera surveys. Deer are born at nearly a 1:1 ratio of does to bucks. Thus, adult sex ratios closer to 1:1 are ideal. In trophy management scenarios adult sex ratio may be skewed towards more bucks, but most deer herds range from 1.5:1 to 3:1.

Important DMAP Reminders for Each Season

- For more information, visit www.mdwfp.com/DMAP and check out the Deer Managers Toolkit.
- If club acreages are incorrect, they should be updated on your DMAP Application. Contact your DMAP Biologist for help updating your DMAP Application.
- Calibrate your body weight scale(s) each year, especially if your club has multiple scales.
- Don't forget to retrieve jawbones from taxidermy bucks. The mature bucks are the most important to have aged.
- Do your best to collect all data completely. Check udders thoroughly for lactation.
- Hunter observation data can provide useful information, especially in clubs with small harvest data sets. Ask your biologist about hunter observation books.